
When Fear Holds Sway

Panic Disorder

Panic disorder is an anxiety disorder characterized by unexpected and repeated episodes of intense fear accompanied by physical symptoms that may include chest pain, heart palpitations, shortness of breath, dizziness, or abdominal distress. These sensations often mimic symptoms of a heart attack or other lifethreatening medical conditions. As a result, the diagnosis of panic disorder is frequently not made until extensive and costly medical procedures fail to provide a correct diagnosis or relief.

Many people with panic disorder develop intense anxiety between episodes. It is not unusual for a person with panic disorder to develop phobias about places or situations where panic attacks have occurred, such as in supermarkets or other everyday situations. As the frequency of panic attacks increases, the person often begins to avoid situations where they fear another attack may occur or where help would not be immediately available. This avoidance may eventually develop into agoraphobia, an inability to go beyond known and safe surroundings because of intense fear and anxiety.



Facts About Panic Disorder

- Approximately 2.4 million American adults ages 18 to 54, or about 1.7 percent of people in this age group in a given year, have panic disorder.¹
- Panic disorder typically develops in late adolescence or early adulthood and is twice as common in women as in men.²
- Panic disorder may coexist with other disorders, most often depression

and substance abuse.³ Appropriate diagnosis and treatment of other disorders are important to successfully treating panic disorder.

Fortunately, research—including studies supported by NIMH—has led to the development of treatments to help people with panic disorder.

Treatments for Panic Disorder

Treatments for panic disorder include medications, commonly the selective serotonin reuptake inhibitors, and a type of psychotherapy known as cognitive-behavioral therapy, which teaches people how to view panic attacks differently and demonstrates ways to reduce anxiety. NIMH is conducting a large-scale study to evaluate the effectiveness of combining these treatments. Appropriate treatment by an experienced professional can reduce or prevent panic attacks in 70 to 90 percent of people with panic disorder. Most patients show significant progress after a few weeks of therapy. Relapses may occur, but they can often be effectively treated just like the initial episode.

Research Findings

Heredity, other biological factors, stressful life events, and thinking in a way that exaggerates relatively normal bodily reactions are all believed to play a role in the onset of panic disorder. The exact cause or causes of panic disorder are unknown and are the subject of intense scientific investigation.

Studies in animals and humans have focused on pinpointing the specific brain areas and circuits involved in anxiety and fear, which underlie anxiety disorders, such as panic disorder. Fear, an emotion that evolved to deal with danger, causes an automatic, rapid protective response that occurs without the need for conscious thought. It has been found that the body's fear response is coordinated by a small structure deep inside the brain, called the amygdala.

The amygdala, although relatively small, is a very complicated structure, and recent research suggests that anxiety disorders may be associated with abnormal activation in the amygdala. One aim of research is to use such basic scientific knowledge to develop new therapies.

For More Information

National Institute of Mental Health (NIMH)

Office of Communications and Public Liaison

Public Inquiries: (301) 443-4513 Media Inquiries: (301) 443-4536 E-mail: nimhinfo@nih.gov Web site: http://www.nimh.nih.gov

All material in this fact sheet is in the public domain and may be copied or reproduced without permission from the Institute. Citation of the source is appreciated.

References

Narrow WE, Rae DS, Regier DA. NIMH epidemiology note: prevalence of anxiety disorders. One-year prevalence best estimates calculated from ECA and NCS data. Population estimates based on U.S. Census estimated residential population age 18 to 54 on July 1, 1998. Unpublished.

²Robins LN, Regier DA, eds. *Psychiatric disorders in America: the Epidemiologic Catchment Area Study.* New York: The Free Press. 1991.

³Regier DA, Rae DS, Narrow WE, et al. Prevalence of anxiety disorders and their comorbidity with mood and addictive disorders. *British Journal of Psychiatry Supplement*, 1998; (34): 24-8.

*Hyman SE, Rudorfer MV. Anxiety disorders. In: Dale DC, Federman DD, eds. *Scientific American® Medicine. Volume 3.* New York: Healtheon/WebMD Corp., 2000, Sect. 13, Subsect. VIII, p. 1.

LeDoux J. Fear and the brain: where have we been, and where are we going? *Biological Psychiatry*, 1998; 44(12): 1229-38.



This is the electronic version of a National Institute of Mental Health (NIMH) publication, available from http://www.nimh.nih.gov/publicat/index.cfm. To order a print copy, call the NIMH Information Center at 301-443-4513 or 1-866-615-6464 (toll-free). Visit the NIMH Web site (http://www.nimh.nih.gov) for information that supplements this publication.

To learn more about NIMH programs and publications, contact the following:

Web address:

http://www.nimh.nih.gov

Phone numbers: 301-443-4513 (local) 1-866-615-6464 (toll-free) 301-443-3431 (TTY)

Street address:
National Institute of Mental Health
Office of Communications
Room 8184, MSC 9663
6001 Executive Boulevard
Bethesda, Maryland 20892-9663 USA

E-mail:

nimhinfo@nih.gov

Fax numbers: 301-443-4279 301-443-5158 (FAX 4U)

This information is in the public domain and can be copied or reproduced without permission from NIMH. To reference this material, we suggest the following format:

National Institute of Mental Health. Title. Bethesda (MD): National Institute of Mental Health, National Institutes of Health, US Department of Health and Human Services; Year of Publication/Printing [Date of Update/Revision; Date of Citation]. Extent. (NIH Publication No XXX XXXX). Availability.

A specific example is:

National Institute of Mental Health. Childhood-Onset Schizophrenia: An Update from the National Institute of Mental Health. Bethesda (MD): National Institute of Mental Health, National Institutes of Health, US Department of Health and Human Services; 2003 [cited 2004 February 24]. (NIH Publication Number: NIH 5124). 4 pages. Available from: http://www.nimh.nih.gov/publicat/schizkids.cfm